



# Non-Alcoholic Fatty Liver Disease

## The Liver

The liver is the largest solid internal organ and it is located underneath the ribcage in the right upper part of the abdomen. Although liver size depends on a person's age, body size and shape, gender, and disease state, in most adults, it is about the size of a football. The liver has many important functions. It acts as a filter for the blood. It metabolizes nutrients and other substances such as medications. It stores energy. It synthesizes proteins that are essential for our body to function, including those that help the blood to clot when we bleed. Although the liver is a very resilient organ that has the ability to repair itself, it is susceptible to damage from many different sources, including viruses, toxins, inherited conditions, and even our body's own immune system.

## Non-Alcoholic Fatty Liver Disease

Non-alcoholic fatty liver disease (NAFLD) is a condition in which fat accumulates in the cells of the liver (hepatocytes). It affects more than 20-30% of adults in North America and is becoming increasingly common. Many people with this condition have no symptoms and medical professionals diagnose them as a part of an investigation into the cause of other abnormal lab tests or after they have had imaging of their abdomen for unrelated reasons. Over many years, excessive fat in the liver may result in significant damage. To help those with the condition, health care professionals must identify those at risk and counsel them on ways to address this disease before significant liver damage occurs. Non-alcoholic fatty liver disease is a treatable condition that individuals can often address with lifestyle modifications.

## Development of Fatty Liver

Non-alcoholic fatty liver disease is not a transmissible condition (i.e., it does not spread from person to person). Rather, it is a condition that occurs most often in individuals who have one or more of the following risk factors for developing the disease:

- obesity,
- diabetes or pre-diabetes,
- high cholesterol,
- high blood pressure,
- sedentary lifestyle, and
- the use of certain medications.

The medical community does not yet know exactly why non-alcoholic fatty liver disease affects some people with the above conditions and not others. It is also incompletely understood why, for some people, fat will accumulate in the liver but not cause significant damage (benign steatosis), whereas in others, it results in significant damage (non-alcoholic steatohepatitis).

As the prevalence of non-alcoholic fatty liver disease is very high in Canada, if you have one or more risk factors for the disease, it may be reasonable to discuss with your health care provider whether you are a candidate for further testing. However, at this time, there are no firm accepted guidelines as to who needs further investigation for this condition, other than those who have unexplained liver enzyme abnormalities.

## Symptoms

Many people who have non-alcoholic fatty liver disease do not have any symptoms but, for those who do, symptoms are generally nonspecific, such as mild fatigue or discomfort in the abdomen. Over many years, the presence of fat may trigger inflammation in the liver, which may lead to formation of scar tissue. If very advanced, the amount of scar tissue in the liver may reach a level termed cirrhosis, which refers to a specific pattern and degree of scar tissue in the liver. For patients with cirrhosis, ongoing damage to the liver may eventually result in signs and symptoms, such as worsening fatigue, fluid accumulation in the abdomen (ascites), bleeding from veins in the esophagus or stomach (varices), and confusion (encephalopathy). The goal in identifying non-alcoholic fatty liver disease is to ensure appropriate monitoring, to offer guidance on how to treat the condition, and to decrease the chances of complications.

---

## Diagnosis

Liver biopsy, which involves the use of a needle to take a sample of liver tissue so that it may be examined under the microscope, is the gold standard for diagnosing non-alcoholic fatty liver disease. While it is currently the only way to confirm the diagnosis, in practice, a combination of blood tests and imaging is usually sufficient for a physician to establish the diagnosis in many patients.

## Investigations

When a person is newly diagnosed with non-alcoholic fatty liver disease, it is important that he or she sees a health care provider with specialized knowledge in that area. This may be a nurse, family doctor, or specialist (hepatologist or gastroenterologist), who will conduct initial investigations to ensure that the diagnosis is certain and to rule out other liver diseases that may co-exist. Most of this information can be obtained through physical examination, blood tests, and imaging of the abdomen.

The role of liver biopsy in the diagnosis of non-alcoholic fatty liver disease remains important in certain circumstances, in particular when the diagnosis is in question. In addition, biopsy can also determine how much, if any, scar tissue is present in the liver. Staging is the process of determining the amount of scar tissue (fibrosis) in the liver and, in the past, liver biopsy was the only tool used. A normal liver with no scar tissue present is stage 0. A combination of increasing amounts of scar tissue in the liver and a change in its pattern increases the classification to a higher stage. Stage 1 is minimal scar tissue and stage 4 is cirrhosis. Although biopsy is a safe procedure, it is an invasive one, and thus does carry some risks, including bleeding and post-procedure pain. One of the drawbacks of liver biopsy, other than its invasive nature, is the fact that it samples only a very small portion of a large organ, and thus is susceptible to sampling error.

## Non-invasive Tools to Measure Fibrosis

Liver biopsy remains the gold standard for the staging of liver disease and it is still a good option for many patients. However, health care providers are increasingly using other effective tools to determine the degree of fibrosis in the liver. Of the emerging alternative staging methods, the following are the most commonly used in Canada.

**FibroScan®** is a non-invasive tool used to assess the degree of fibrosis in the liver. It is a technique used to measure liver stiffness, which is closely related to the degree of fibrosis in the liver. The scan involves the painless placement of a probe on the surface of the skin, and takes only a few minutes to complete. The area sampled is approximately 100 times that seen on a typical liver biopsy. This procedure results in a reliable reading for most people. FibroScan® is available at a number of centres across Canada.

**FibroTest and APRI** (AST-to-Platelet Ratio Index) are non-invasive tools that rely on calculations based on blood tests to measure the degree of fibrosis in the liver. FibroTest is likely not covered by medical plans and therefore will have associated out-

of-pocket expenses. APRI is derived from a calculation using a simple equation based on common blood tests.

Regardless of the tool used to estimate fibrosis, expert interpretation is essential in ensuring the information gained is useful in making treatment decisions. In addition, liver biopsy may still provide valuable information regarding disease activity that non-invasive tools cannot. Your health care provider will be able to determine which tests are best suited to your situation.

## Management

Not everyone who has non-alcoholic fatty liver disease is at risk of developing progressive liver disease. However, because the management of this condition focuses on healthy lifestyle that may also decrease the chances of developing cardiovascular disease, most patients will benefit in the long term from discussing the management of this condition with their health care provider.

### Alcohol

Significant alcohol consumption is a known risk factor for accumulating fat in the liver. Most consider significant alcohol consumption to be more than 2 drinks a day. In someone with established non-alcoholic fatty liver disease, limiting intake to no more than 1-2 drinks per day (one drink is 5oz of wine, 1.5oz of spirits, or 355mL of beer) and not consuming alcohol every day is a reasonable goal. In fact, complete abstinence from alcohol may be preferable.

### Other Medical Conditions

Other medical conditions, such as high cholesterol, diabetes, and high blood pressure, increase an individual's risk of developing non-alcoholic fatty liver disease, and may worsen its severity if already present. Thus, optimal control of these risk factors is central to managing non-alcoholic fatty liver disease. In general, the medications typically used to treat these risk factors are also safe in people with liver disease, and patients should discuss this further with their health care providers.

### Medications

Researchers have studied many medications, vitamins, and supplements for their potential beneficial effects in the management of non-alcoholic fatty liver disease. Unfortunately, to date, none of them have found evidence to justify their universal and widespread use for individuals with this condition. Some practitioners continue to use Vitamin E and pioglitazone (Actos®; a medication used to treat diabetes), in certain situations, but there are limitations with respect to both efficacy and safety that patients should discuss in detail with their health care providers before pursuing treatment with these agents.

### Lifestyle Changes

While research is ongoing to identify medications suitable for widespread use in the management of non-alcoholic fatty liver disease, multiple well-designed trials have demonstrated lifestyle changes (exercise and dietary) that result in weight loss

to be an effective way to manage this disease.

Individuals who are able to achieve a weight loss of between 3-10% of their total body weight by increasing activity levels and improving diet also achieve improvement in liver enzymes, reduced amounts of fat in their liver, and reduced inflammation in the liver.

The current *Canadian Physical Activity Guidelines* advise that all adults 18-64 years old should accumulate at least 150 minutes of moderate- to vigorous-intensity aerobic activity a week, in bouts of 10 minutes or more, to achieve health benefits. This is also a reasonable goal for people with non-alcoholic fatty liver disease who are not already exercising at this level. Dietary changes that will be of benefit include limiting intake of high fat and high sugar foods, limiting portion size, and eating regular balanced meals by following *Canada's Food Guide*. Your health care provider can give you further advice on optimizing your exercise regimen and diet. Following the advice of a personal trainer and/or dietitian may also help you reach your goals.

## Outlook

The future of non-alcoholic fatty liver disease lies in the education of health care providers and the public so that practitioners can identify and care for the many people with this condition. We hope that research into new therapeutic options for non-alcoholic fatty liver disease and an ongoing interest in learning more about this condition will, over the coming years, result in the following:

- an improved understanding of the pathophysiology of this condition,
- safe and effective medications for widespread use, and
- improved tools to diagnose and monitor this condition.

## NOTES:

---

---

---

---

---

---

---

---

---

---



## ABOUT US

As the Canadian leader in providing trusted, evidence-based information on all areas of the gastrointestinal (GI) tract, the Gastrointestinal Society is committed to improving the lives of people with GI and liver conditions, supporting research, advocating for appropriate patient access to healthcare, and promoting gastrointestinal and liver health.

The *Inside Tract*® newsletter provides the latest news on GI research, disease and disorder treatments (e.g., medications, nutrition), and a whole lot more. If you have any kind of digestive problem, then you'll want this timely, informative publication.

### Please subscribe today!

The GI Society, in partnership with the Canadian Society of Intestinal Research, produced this pamphlet under the guidance of affiliated healthcare professionals. This document is not intended to replace the knowledge, diagnosis, or care of your physician.

© Gastrointestinal Society 2013

## SUBSCRIPTION/DONATION FORM

### Purchase Information

- I enclose a donation of \$ \_\_\_\_\_ (Eligible for a tax receipt)  
 I enclose a \$20 annual subscription fee (\$30 outside Canada)

Total \$ \_\_\_\_\_

Name (Mr./Mrs./Ms./Dr./Other \_\_\_\_\_ Please circle one)

Company Name (Optional)

Street Address

City, Province, Postal Code

Daytime Phone \_\_\_\_\_ Email \_\_\_\_\_

- Patient     Friend/Family Member     Professional

**Diagnosis/Area of Interest** (This optional information helps us to determine which topics might be of interest to you and is kept strictly confidential. We never sell our lists.)

Would you like more information? Please list your topics.

**Payment:**  Visa     Mastercard     Cheque Enclosed\*

Credit Card Number \_\_\_\_\_ Expiry Date \_\_\_\_\_

\*Make cheque payable to the **GI Society**.

**Gastrointestinal Society**  
855 West 12th Avenue  
Vancouver, BC V5Z 1M9  
Phone: 604-875-4875

**Toll-free:** 1-866-600-4875  
**Fax:** 604-875-4429  
**Email:** info@badgut.org  
**Website:** www.badgut.org

**Charitable Registration Number:** 817065352RR0001